

TOP LOCAL COHOMOLOGY MODULES WITH SPECIFIED ATTACHED PRIMES

MOHAMMAD T. DIBAEI AND RAHELEH JAFARI

ABSTRACT. Let $(R, \mathfrak{m})$ be a complete Noetherian local ring and let M be a finite R -module of positive Krull dimension n . It is shown that any subset T of $\text{Assh}_R(M)$ can be expressed as the set of attached primes of the top local cohomology module $H_{\mathfrak{a}}^n(M)$ for some ideal $\mathfrak{a}$ of R . Moreover if $\mathfrak{a}$ is an ideal of R such that the set of attached primes of $H_{\mathfrak{a}}^n(M)$ is a non-empty proper subset of $\text{Assh}_R(M)$, then $H_{\mathfrak{a}}^n(M) \cong H_{\mathfrak{b}}^n(M)$ for some ideal $\mathfrak{b}$ of R with $\dim_R(R/\mathfrak{b}) = 1$.

1. INTRODUCTION

Throughout $(R, \mathfrak{m})$ is a commutative Noetherian local ring with maximal ideal $\mathfrak{m}$, M is a non-zero finite (i.e. finitely generated) R -module with positive Krull dimension $n := \dim_R(M)$ and $\mathfrak{a}$ denotes an ideal of R . Recall that for an R -module N , a prime ideal $\mathfrak{p}$ of R is said to be an *attached prime* of N , if $\mathfrak{p} = \text{Ann}_R(N/K)$ for some submodule K of N (see [5]). The set of attached primes of N is denoted by $\text{Att}_R(N)$. If N is an Artinian R -module so that N admits a reduced secondary representation $N = N_1 + \cdots + N_r$ such that N_i is $\mathfrak{p}_i$ -secondary, $i = 1, \dots, r$, then $\text{Att}_R(N) = \{\mathfrak{p}_1, \dots, \mathfrak{p}_r\}$ is a finite set.

Denote by $H_{\mathfrak{a}}^n(M)$ the n th right derived functor of

$$\Gamma_{\mathfrak{a}}(M) = \{x \in M \mid \mathfrak{a}^r x = 0 \text{ for some positive integer } r\}$$

applied to M . It is well-known that $H_{\mathfrak{a}}^n(M)$ is an Artinian module. Macdonald and Sharp, in [5], studied $H_{\mathfrak{m}}^n(M)$ and showed that $\text{Att}_R(H_{\mathfrak{m}}^n(M)) = \text{Assh}_R(M)$ where $\text{Assh}_R(M) := \{\mathfrak{p} \in \text{Ass}_R(M) \mid \dim_R(R/\mathfrak{p}) = n\}$. It is shown in [4, Theorem A], that for any arbitrary ideal $\mathfrak{a}$ of R , $\text{Att}_R(H_{\mathfrak{a}}^n(M)) = \{\mathfrak{p} \in \text{Ass}_R(M) \mid H_{\mathfrak{a}}^n(R/\mathfrak{p}) \neq 0\}$ which is a subset of $\text{Assh}_R(M)$. In [3], the structure of $H_{\mathfrak{a}}^n(M)$ is studied by the first author and Yassemi and they showed that, in case R is complete, for any pair of ideals $\mathfrak{a}$ and $\mathfrak{b}$ of R , if $\text{Att}_R(H_{\mathfrak{a}}^n(M)) = \text{Att}_R(H_{\mathfrak{b}}^n(M))$, then $H_{\mathfrak{a}}^n(M) \cong H_{\mathfrak{b}}^n(M)$. They also raised the following question in [2, Question 2.9] which is the main object of this paper.

Question. For any subset T of $\text{Assh}_R(M)$, is there an ideal $\mathfrak{a}$ of R such that $\text{Att}_R(H_{\mathfrak{a}}^n(M)) = T$?

This paper provides a positive answer for this question in the case R is complete.

2. MAIN RESULT

In this section we assume that R is complete with respect to the $\mathfrak{m}$ -adic topology. As mentioned above, $\text{Att}_R(H_{\mathfrak{m}}^n(M)) = \text{Assh}_R(M)$ and $\text{Att}_R(H_R^n(M)) = \emptyset$ is the empty set. Also $\text{Att}_R(H_{\mathfrak{a}}^n(M)) \subseteq \text{Assh}_R(M)$ for all ideals $\mathfrak{a}$ of R . Our aim is to show that as $\mathfrak{a}$ varies over ideals of R , the set $\text{Att}_R(H_{\mathfrak{a}}^n(M))$ takes all possible subsets of $\text{Assh}_R(M)$ (see Theorem 2.8). In the following results we always assume that T is a non-empty proper subset of $\text{Assh}_R(M)$.

In our first result we find a characterization for a subset of $\text{Assh}_R(M)$ to be the set of attached primes of the top local cohomology of M with respect to an ideal $\mathfrak{a}$.

2000 *Mathematics Subject Classification.* 13D45, 13D07.

Key words and phrases. Local cohomology, Associated primes, Attached primes

The research of the first author was in part supported from IPM (No. 00000000).

Proposition 2.1. *Assume that $n := \dim_R(M) \geq 1$ and that T is a proper non-empty subset of $\text{Assh}_R(M)$. Set $\text{Assh}_R(M) \setminus T = \{\mathfrak{q}_1, \dots, \mathfrak{q}_r\}$. The following statements are equivalent.*

- (i) *There exists an ideal $\mathfrak{a}$ of R such that $\text{Att}_R(H_{\mathfrak{a}}^n(M)) = T$.*
- (ii) *For each i , $1 \leq i \leq r$, there exists $Q_i \in \text{Supp}_R(M)$ with $\dim_R(R/Q_i) = 1$ such that*

$$\bigcap_{\mathfrak{p} \in T} \mathfrak{p} \not\subseteq Q_i \quad \text{and} \quad \mathfrak{q}_i \subseteq Q_i.$$

With Q_i , $1 \leq i \leq r$, as above, $\text{Att}_R(H_{\mathfrak{a}}^n(M)) = T$ where $\mathfrak{a} = \bigcap_{i=1}^r Q_i$.

Proof. (i) $\Rightarrow$ (ii). By [4, Theorem A], $H_{\mathfrak{a}}^n(R/\mathfrak{p}) \neq 0$ for all $\mathfrak{p} \in T$, that is $\mathfrak{a} + \mathfrak{p}$ is $\mathfrak{m}$ -primary for all $\mathfrak{p} \in T$ (by Lichtenbaum-Hartshorne Theorem). On the other hand, for $1 \leq i \leq r$, $\mathfrak{q}_i \not\subseteq T$ which is equivalent to say that $\mathfrak{a} + \mathfrak{q}_i$ is not an $\mathfrak{m}$ -primary ideal. Hence there exists a prime ideal $Q_i \in \text{Supp}_R(M)$ such that $\dim_R(R/Q_i) = 1$ and $\mathfrak{a} + \mathfrak{q}_i \subseteq Q_i$. It follows that $\bigcap_{\mathfrak{p} \in T} \mathfrak{p} \not\subseteq Q_i$.

(ii) $\Rightarrow$ (i). Set $\mathfrak{a} := \bigcap_{i=1}^r Q_i$. For each i , $1 \leq i \leq r$, $\mathfrak{a} + \mathfrak{q}_i \subseteq Q_i$ implies that $\mathfrak{a} + \mathfrak{q}_i$ is not $\mathfrak{m}$ -primary and so $H_{\mathfrak{a}}^n(R/\mathfrak{q}_i) = 0$. Thus $\text{Att}_R H_{\mathfrak{a}}^n(M) \subseteq T$. Assume $\mathfrak{p} \in T$ and $Q \in \text{Supp}(M)$ such that $\mathfrak{a} + \mathfrak{p} \subseteq Q$. Then $Q_i \subseteq Q$ for some i , $1 \leq i \leq r$. Since $\mathfrak{p} \not\subseteq Q_i$, we have $Q_i \neq Q$, so $Q = \mathfrak{m}$. Hence $\mathfrak{a} + \mathfrak{p}$ is $\mathfrak{m}$ -primary ideal. Now, by Lichtenbaum-Hartshorne Theorem, and by [4, Theorem A], it follows that $\mathfrak{p} \in \text{Att}_R(H_{\mathfrak{a}}^n(M))$. $\square$

Corollary 2.2. *If $H_{\mathfrak{a}}^n(M) \neq 0$ then there is an ideal $\mathfrak{b}$ of R such that $\dim_R(R/\mathfrak{b}) \leq 1$ and $H_{\mathfrak{a}}^n(M) \cong H_{\mathfrak{b}}^n(M)$.*

Proof. If $\text{Att}_R(H_{\mathfrak{a}}^n(M)) = \text{Assh}_R(M)$, then $H_{\mathfrak{a}}^n(M) = H_{\mathfrak{m}}^n(M)$. Otherwise $n \geq 1$ and $\text{Att}_R(H_{\mathfrak{a}}^n(M))$ is a proper subset of $\text{Assh}_R(M)$. Set $\text{Assh}_R(M) \setminus \text{Att}_R(H_{\mathfrak{a}}^n(M)) := \{\mathfrak{q}_1, \dots, \mathfrak{q}_r\}$. By Proposition 2.1, there are $Q_i \in \text{Supp}_R(M)$ with $\dim_R(R/Q_i) = 1$, $i = 1, \dots, r$, such that $\text{Att}_R(H_{\mathfrak{a}}^n(M)) = \text{Att}_R(H_{\mathfrak{b}}^n(M))$ with $\mathfrak{b} = \bigcap_{i=1}^r Q_i$. Now, by [3, Theorem 1.6], we have $H_{\mathfrak{a}}^n(M) \cong H_{\mathfrak{b}}^n(M)$. As $\dim(R/\mathfrak{b}) = 1$, the proof is complete. $\square$

Corollary 2.3. *If $\dim_R(M) = 1$ then any subset T of $\text{Assh}_R(M)$ is equal to the set $\text{Att}_R(H_{\mathfrak{a}}^1(M))$ for some ideal $\mathfrak{a}$ of R .*

Proof. With notations as in Proposition 2.1, we take $Q_i = \mathfrak{q}_i$ for $i = 1, \dots, r$. $\square$

By a straightforward argument one may notice that the condition “complete” is superficial, for if T is a non-empty proper subset of $\text{Assh}_R(M)$, then $T = \text{Att}_R(H_{\mathfrak{a}}^1(M))$, where $\mathfrak{a} = \bigcap_{\mathfrak{p} \in \text{Assh}_R(M) \setminus T} \mathfrak{p}$.

The following is an example to Proposition 2.1.

Example 2.4. *Set $R = k[[X, Y, Z, W]]$, where k is a field and X, Y, Z, W are independent indeterminates. Then R is a complete Noetherian local ring with maximal ideal $\mathfrak{m} = (X, Y, Z, W)$. Consider prime ideals*

$$\mathfrak{p}_1 = (X, Y) \quad , \quad \mathfrak{p}_2 = (Z, W) \quad , \quad \mathfrak{p}_3 = (Y, Z) \quad , \quad \mathfrak{p}_4 = (X, W)$$

and set $M = \frac{R}{\mathfrak{p}_1 \mathfrak{p}_2 \mathfrak{p}_3 \mathfrak{p}_4}$ as an R -module, so that we have $\text{Assh}_R(M) = \{\mathfrak{p}_1, \mathfrak{p}_2, \mathfrak{p}_3, \mathfrak{p}_4\}$ and $\dim_R(M) = 2$. We get $\{\mathfrak{p}_i\} = \text{Att}_R(H_{\mathfrak{a}_i}^2(M))$, where $\mathfrak{a}_1 = \mathfrak{p}_2, \mathfrak{a}_2 = \mathfrak{p}_1, \mathfrak{a}_3 = \mathfrak{p}_4, \mathfrak{a}_4 = \mathfrak{p}_3$, and

$\{\mathfrak{p}_i, \mathfrak{p}_j\} = \text{Att}_R(H_{\mathfrak{a}_{ij}}^2(M))$, where

$$\begin{aligned}\mathfrak{a}_{12} &= (Y^2 + YZ, Z^2 + YZ, X^2 + XW, W^2 + WX), \\ \mathfrak{a}_{34} &= (Z^2 + ZW, X^2 + YX, Y^2 + YX, W^2 + WZ), \\ \mathfrak{a}_{13} &= (Z^2 + XZ, W^2 + WY, X^2 + XZ), \\ \mathfrak{a}_{14} &= (W^2 + WY, Z^2 + ZY, Y^2 + YW), \\ \mathfrak{a}_{23} &= (X^2 + XZ, Y^2 + WY, W^2 + ZW), \\ \mathfrak{a}_{24} &= (X^2 + XZ, Y^2 + WY, Z^2 + ZW).\end{aligned}$$

Finally, we have $\{\mathfrak{p}_i, \mathfrak{p}_j, \mathfrak{p}_k\} = \text{Att}_R(H_{\mathfrak{a}_{ijk}}^2(M))$, where $\mathfrak{a}_{123} = (X, W, Y + Z)$, $\mathfrak{a}_{234} = (X, Y, W + Z)$, $\mathfrak{a}_{134} = (Z, W, Y + X)$.

Lemma 2.5. Assume that $n := \dim_R(M) \geq 2$, and that T is a non-empty subset of $\text{Assh}_R(M)$ such that $\bigcap_{\mathfrak{p} \in T} \mathfrak{p} \not\subseteq \bigcap_{\mathfrak{q} \in \text{Assh}_R(R/\sum_{\mathfrak{p} \in T'} \mathfrak{p})} \mathfrak{q}$, where $T' = \text{Assh}_R(M) \setminus T$. Then there exists a prime ideal

$Q \in \text{Supp}_R(M)$ with $\dim_R(R/Q) = 1$ and $\text{Att}_R(H_Q^n(M)) = T$.

Proof. Set $s := \text{ht}_M(\sum_{\mathfrak{p} \in T'} \mathfrak{p})$. We have $s \leq n - 1$, otherwise $\text{Assh}_R(R/\sum_{\mathfrak{p} \in T'} \mathfrak{p}) = \{\mathfrak{m}\}$ which contradicts the condition $\bigcap_{\mathfrak{p} \in T} \mathfrak{p} \not\subseteq \bigcap_{\mathfrak{q} \in \text{Assh}_R(R/\sum_{\mathfrak{p} \in T'} \mathfrak{p})} \mathfrak{q}$. As R is catenary, we have $\dim_R(R/\sum_{\mathfrak{p} \in T'} \mathfrak{p}) = n - s$.

We first prove, by induction on j , $0 \leq j \leq n - s - 1$, that there exists a chain of prime ideals $Q_0 \subset Q_1 \subset \cdots \subset Q_j \subset \mathfrak{m}$ such that $Q_0 \in \text{Assh}_R(R/\sum_{\mathfrak{p} \in T'} \mathfrak{p})$, $\dim_R(R/Q_j) = n - s - j$ and $\bigcap_{\mathfrak{p} \in T} \mathfrak{p} \not\subseteq Q_j$. There is $Q_0 \in \text{Assh}_R(R/\sum_{\mathfrak{p} \in T'} \mathfrak{p})$ such that $\bigcap_{\mathfrak{p} \in T} \mathfrak{p} \not\subseteq Q_0$. Note that $\dim_R(R/Q_0) = \dim_R(R/\sum_{\mathfrak{p} \in T'} \mathfrak{p}) = n - s$. Now, assume that $0 < j \leq n - s - 1$ and that we have proved the existence of a chain $Q_0 \subset Q_1 \subset \cdots \subset Q_{j-1}$ of prime ideals such that $Q_0 \in \text{Assh}_R(R/\sum_{\mathfrak{p} \in T'} \mathfrak{p})$, $\dim_R(R/Q_j) = n - s - (j - 1)$ and that $\bigcap_{\mathfrak{p} \in T} \mathfrak{p} \not\subseteq Q_{j-1}$. Note that we have $n - s - (j - 1) = n - s + 1 - j \geq 2$. Therefore the set V defined as

$$V = \{\mathfrak{q} \in \text{Supp}_R(M) \mid Q_{j-1} \subset \mathfrak{q} \subset \mathfrak{q}' \subseteq \mathfrak{m}, \dim_R(R/\mathfrak{q}) = n - s - j, \mathfrak{q}' \in \text{Spec}(R) \text{ and } \dim_R(R/\mathfrak{q}') = n - s - j - 1\}$$

is non-empty and so, by Ratliff's weak existence theorem [6, Theorem 31.2], is not finite. As $\bigcap_{\mathfrak{p} \in T} \mathfrak{p} \not\subseteq Q_{j-1}$, we have $Q_{j-1} \subset Q_{j-1} + \bigcap_{\mathfrak{p} \in T} \mathfrak{p}$. If, for $\mathfrak{q} \in V$, $\bigcap_{\mathfrak{p} \in T} \mathfrak{p} \subseteq \mathfrak{q}$, then $\mathfrak{q}$ is a minimal prime of $Q_{j-1} + \bigcap_{\mathfrak{p} \in T} \mathfrak{p}$. As V is an infinite set, there is $Q_j \in V$ such that $\bigcap_{\mathfrak{p} \in T} \mathfrak{p} \not\subseteq Q_j$. Thus the induction is complete. Now by taking $Q := Q_{n-s-1}$ and by Proposition 2.1, the claim follows. $\square$

Corollary 2.6. Assume that $n := \dim_R(M) \geq 2$ and T is a non-empty subset of $\text{Assh}_R(M)$ with $|T| = |\text{Assh}_R(M)| - 1$. Then there is an ideal $\mathfrak{a}$ of R such that $\text{Att}_R(H_{\mathfrak{a}}^n(M)) = T$.

Proof. Note that $\text{Assh}_R(M) \setminus T$ is a singleton set $\{\mathfrak{q}\}$, say, and so $\text{ht}_M(\mathfrak{q}) = 0$ and $\bigcap_{\mathfrak{p} \in T} \mathfrak{p} \not\subseteq \mathfrak{q}$. Therefore, by Lemma 2.5, the result follows. $\square$

Lemma 2.7. Assume that $n := \dim_R(M) \geq 2$ and $\mathfrak{a}_1$ and $\mathfrak{a}_2$ are ideals of R . Then there exists an ideal $\mathfrak{b}$ of R such that $\text{Att}_R(H_{\mathfrak{b}}^n(M)) = \text{Att}_R(H_{\mathfrak{a}_1}^n(M)) \cap \text{Att}_R(H_{\mathfrak{a}_2}^n(M))$.

Proof. Set $T_1 = \text{Att}_R(H_{\mathfrak{a}_1}^n(M))$ and $T_2 = \text{Att}_R(H_{\mathfrak{a}_2}^n(M))$. We may assume that $T_1 \cap T_2$ is a non-empty proper subset of $\text{Assh}_R(M)$. Assume that $\mathfrak{q} \in \text{Assh}_R(M) \setminus (T_1 \cap T_2) = (\text{Assh}_R(M) \setminus$

$T_1) \cup (\text{Assh}_R(M) \setminus T_2)$. By Proposition 2.1, there exists $Q \in \text{Supp}_R(M)$ with $\dim_R(R/Q) = 1$ such that $\mathfrak{q} \subseteq Q$ and $\bigcap_{\mathfrak{p} \in T_1 \cap T_2} \mathfrak{p} \not\subseteq Q$. Now, by Proposition 2.1, again there exists an ideal $\mathfrak{b}$ of R such that $\text{Att}_R(H_{\mathfrak{b}}^n(M)) = T_1 \cap T_2$. $\square$

Now we are ready to present our main result.

Theorem 2.8. *Assume that $T \subseteq \text{Assh}_R(M)$, then there exists an ideal $\mathfrak{a}$ of R such that $T = \text{Att}_R(H_{\mathfrak{a}}^n(M))$.*

Proof. By Corollary 2.3, we may assume that $\dim_R(M) \geq 2$ and that T is a non-empty proper subset of $\text{Assh}_R(M)$. Set $T = \{\mathfrak{p}_1, \dots, \mathfrak{p}_t\}$ and $\text{Assh}_R(M) \setminus T = \{\mathfrak{p}_{t+1}, \dots, \mathfrak{p}_{t+r}\}$. We use induction on r . For $r = 1$, Corollary 2.6 proves the first step of induction. Assume that $r > 1$ and that the case $r - 1$ is proved. Set $T_1 = \{\mathfrak{p}_1, \dots, \mathfrak{p}_t, \mathfrak{p}_{t+1}\}$ and $T_2 = \{\mathfrak{p}_1, \dots, \mathfrak{p}_t, \mathfrak{p}_{t+2}\}$. By induction assumption there exist ideals $\mathfrak{a}_1$ and $\mathfrak{a}_2$ of R such that $T_1 = \text{Att}_R(H_{\mathfrak{a}_1}^n(M))$ and $T_2 = \text{Att}_R(H_{\mathfrak{a}_2}^n(M))$. Now by the Lemma 2.7 there exists an ideal $\mathfrak{a}$ of R such that $T = T_1 \cap T_2 = \text{Att}_R(H_{\mathfrak{a}}^n(M))$. $\square$

Corollary 2.9. *(See [1, Corollary 1.7]) With the notations as in Theorem 2.8, the number of non-isomorphic top local cohomology modules of M with respect to all ideals of R is equal to $2^{|\text{Assh}_R(M)|}$.*

Proof. It follows from Theorem 2.8 and [3, Theorem 1.6]. $\square$

Acknowledgment. The authors would like to thank the referee for her/his comments.

REFERENCES

1. F. W. Call, *On local cohomology modules*, J. Pure Appl. Algebra **43** (1986), no. 2, 111–117.
2. M. T. Dibaei and S. Yassemi, *Some rigidity results for highest order local cohomology modules*, Algebra Colloq., to appear.
3. M. T. Dibaei and S. Yassemi, *Top local cohomology modules*, Algebra Colloq., to appear.
4. M. T. Dibaei and S. Yassemi, *Attached primes of the top local cohomology modules with respect to an ideal*, Arch. Math. (Basel) **84** (2005), no. 4, 292–297.
5. I. G. Macdonald and R. Y. Sharp, *An elementary proof of the non-vanishing of certain local cohomology modules*, Quart. J. Math. Oxford **23** (1972), 197–204.
6. H. Matsumura, *Commutative ring theory*, Cambridge Studies in Advanced Mathematics, 8. Cambridge University Press, Cambridge, 1986.

MOHAMMAD T. DIBAEI, FACULTY OF MATHEMATICAL SCIENCES, TEACHER TRAINING UNIVERSITY, TEHRAN, IRAN,
AND INSTITUTE FOR THEORETICAL PHYSICS AND MATHEMATICS (IPM), TEHRAN, IRAN.

E-mail address: dibaeint@ipm.ir

RAHELEH JAFARI, FACULTY OF MATHEMATICAL SCIENCES, TEACHER TRAINING UNIVERSITY, TEHRAN, IRAN

E-mail address: jafarirahale@yahoo.com